

## METHODS AND COMPOSITIONS FOR TREATING EPILEPSY

### FIELD OF THE INVENTION

[0001] The invention relates to compositions that treat epilepsy and can be  
5 taken as dietary supplements or medication. These compositions contain yeast  
cells obtainable by growth in electromagnetic fields with specific frequencies and  
field strengths.

### BACKGROUND OF THE INVENTION

[0002] Epilepsy is a chronic illness caused by abnormality in the central  
10 nervous system. An epileptic seizure is a brief, excessive surge of electrical  
activity in the brain that causes a change in consciousness, sensation and behavior.  
During an epileptic seizure, the regulatory systems that maintain the normal  
balance between excitation and inhibition of the brain's electrical activity break  
down. There may be a loss of inhibitory nerve cells or an overproduction of an  
15 excitatory neurotransmitter. Groups of abnormal cells are activated synchronously,  
creating a storm of electrical activity.

[0003] Patients taking anticonvulsant drugs display a broad spectrum of  
side-effects. The widely used drug carbamazepine, shows side effects such as  
dizziness, ataxia, drowsiness and reduction of alertness. See, A. Delcker et al.,  
20 Eur. Neuropsychopharmacol., 7, pp. 213-8 (1997). Valproic acid may precipitate  
metabolic disorders, liver disease, gastrointestinal symptomatology, excessive  
bodyweight gain and alopecia. See, S. J. Wallace, Drug Saf, 15, pp. 378-93

(1996). Barbiturates precipitates metabolic bone disease and rash. See, S. J. Wallace, Drug Saf, 15, pp. 378-93 (1996). Therefore, there is a need on the market for anticonvulsant medication with fewer side effects.

### **SUMMARY OF THE INVENTION**

5 [0004] The composition of the invention assists in the recovery of epilepsy and can be taken as dietary supplements in the form of health drinks or pills.

[0005] This invention embraces a composition comprising a plurality of yeast cells that have been cultured in the presence of an alternating electric field having a frequency in the range of about 10200 to 13040 MHz and a field strength  
10 in the range of about 20 to 600 mV/cm. In one embodiment, the frequency is in the range of about 10200-10270, 12330-12390, or 12970-13040 MHz. In another embodiment, the field strength is in the range of about 200-500 mV/cm. The yeast cells are cultured in the alternating electric field for a period of time sufficient to increase the capability of said plurality of yeast cells to have an anti-seizure effect  
15 or treat epilepsy as compared to unactivated yeast cells. In one embodiment, the composition comprising the activated yeast cells reduces the occurrence of epileptic seizures in mammals. Preferably, the mammal is human. In one embodiment, the human has seizure activity. In one embodiment, the frequency and/or the field strength of the alternating electric field can be altered within the  
20 aforementioned ranges during said period of time. In other words, the yeast cells can be exposed to a series of electromagnetic fields. An exemplary period of time is about 140-210 hours.

[0006] Also included in this invention is a composition comprising a plurality of yeast cells that have been cultured under acidic conditions in an  
25 alternating electric field having a frequency in the range of about 12970-13040 MHz and a field strength in the range of about 260 to 510 mV/cm (e.g., 260-280, 330-360, 350-380, 430-470 or 470-510 mV/cm). In one embodiment, the yeast cells are exposed to a series of electromagnetic fields. An exemplary period of time is about 180-210 hours.

[0007] Included in this invention are also methods of making the above compositions and methods of treating a subject with epilepsy.

[0008] Yeast cells that can be included in this composition can all be derived from parent strains publically available from the China General

- 5 Microbiological Culture Collection Center ("CGMCC") (China Committee for Culture Collection of Microorganisms, Institute of Microbiology, Chinese Academy of Sciences, Haidian, P.O. BOX 2714, Beijing, 100080, China). Useful yeast species include, but are not limited to *Schizosaccharomyces pombe*, *Saccharomyces sake*, *Saccharomyces uvarum*, *Saccharomyces rouxii*,  
10 *Saccharomyces carlsbergensis*, *Rhodotorula aurantiaca* and *Saccharomyces cerevisiae*. In one embodiment, the yeast species is *Saccharomyces carlsbergensis* Hansen or *Saccharomyces cerevisiae* Hansen. For instance, the yeast cells can be of the strain *Saccharomyces cerevisiae* Hansen IFFI1335. In one embodiment, the yeast cells are from the strains selected from the group consisting of  
15 *Saccharomyces cerevisiae* Hansen AS2.501, AS2.502, AS2.503, AS2.504, AS2.535, AS2.558, AS2.560, AS2.561, AS2.443 and AS2.562. Other useful yeast species are illustrated in Table 1.

- [0009] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in  
20 the art to which this invention belongs. Exemplary methods and materials are described below, although methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention. All publications and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including  
25 definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting. Throughout this specification and claims, the word "comprise," or variations such as "comprises" or "comprising" will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

- 30 [0010] Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

### **BRIEF DESCRIPTION OF THE DRAWINGS**

[0011] Fig. 1 is a schematic diagram showing an exemplary apparatus for activating yeast cells using electromagnetic fields. 1: yeast culture; 2: container; 3: power supply.

- 5 [0012] Fig. 2 is a schematic diagram showing an exemplary apparatus for making yeast compositions of the invention. The apparatus comprises a signal generator (such as models 83721B and 83741A manufactured by HP) and interconnected containers A, B and C.

### **DETAILED DESCRIPTION OF THE INVENTION**

- 10 [0013] This invention is based on the discovery that certain yeast strains can be activated by electromagnetic fields ("EMF") having specific frequencies and field strengths to produce agents useful in treating epilepsy. Yeast compositions comprising activated yeast cells can be used as medication or dietary supplements in the form of health drinks or dietary pills. In one embodiment, the yeast  
15 compositions of this invention have an anti-seizure effect in a mammal. In one embodiment, the mammal is human.

- [0014] Since the activated yeast cells contained in these yeast compositions have been cultured to endure acidic conditions of pH 2.5-4.2, the compositions are stable in the stomach and can pass on to the intestines. Once in the intestines, the  
20 yeast cells are ruptured by various digestive enzymes, and agents for regulating the central nervous system are released and readily absorbed.

#### **I. Yeast Strains Useful in the Invention**

- [0015] The types of yeasts useful in this invention include, but are not limited to, yeasts of the genera *Saccharomyces*, *Schizosaccharomyces*, and  
25 *Rhodotorula*.

- [0016] Exemplary species within the above-listed genera include, but are not limited to, the species illustrated in Table 1. Yeast strains useful in this invention can be obtained from laboratory cultures, or from publically accessible culture depositories, such as CGMCC and the American Type Culture Collection,

10801 University Boulevard, Manassas, VA 20110-2209. Non-limiting examples of useful strains (with the accession numbers of CGMCC) are illustrated in Table 1. In general, yeast strains preferred in this invention are those used for fermentation in the food and wine industries. As a result, compositions containing these yeast cells are safe for human consumption.

[0017] The preparation of the yeast compositions of this invention is not limited to starting with a pure strain of yeast. A yeast composition of the invention may be produced by culturing a mixture of yeast cells of different species or strains.

Table 1 Exemplary Yeast Strains

<i>Saccharomyces cerevisiae</i> Hansen				
ACCC2034	ACCC2035	ACCC2036	ACCC2037	ACCC2038
ACCC2039	ACCC2040	ACCC2041	ACCC2042	AS2. 1
AS2. 4	AS2. 11	AS2. 14	AS2. 16	AS2. 56
AS2. 69	AS2. 70	AS2. 93	AS2. 98	AS2. 101
AS2. 109	AS2. 110	AS2. 112	AS2. 139	AS2. 173
AS2. 174	AS2. 182	AS2. 196	AS2. 242	AS2. 336
AS2. 346	AS2. 369	AS2. 374	AS2. 375	AS2. 379
AS2. 380	AS2. 382	AS2. 390	AS2. 393	AS2. 395
AS2. 396	AS2. 397	AS2. 398	AS2. 399	AS2. 400
AS2. 406	AS2. 408	AS2. 409	AS2. 413	AS2. 414
AS2. 415	AS2. 416	AS2. 422	AS2. 423	AS2. 430
AS2. 431	AS2. 432	AS2. 451	AS2. 452	AS2. 453
AS2. 458	AS2. 460	AS2. 463	AS2. 467	AS2. 486
AS2. 501	AS2. 502	AS2. 503	AS2. 504	AS2. 516
AS2. 535	AS2. 536	AS2. 558	AS2. 560	AS2. 561
AS2. 562	AS2. 576	AS2. 593	AS2. 594	AS2. 614
AS2. 620	AS2. 628	AS2. 631	AS2. 666	AS2. 982
AS2. 1190	AS2. 1364	AS2. 1396	IFFI1001	IFFI1002
IFFI1005	IFFI1006	IFFI1008	IFFI1009	IFFI1010

5	IFFI1012	IFFI1021	IFFI1027	IFFI1037	IFFI1042
	IFFI1043	IFFI1045	IFFI1048	IFFI1049	IFFI1050
	IFFI1052	IFFI1059	IFFI1060	IFFI1062	IFFI1063
	IFFI1202	IFFI1203	IFFI1206	IFFI1209	IFFI1210
	IFFI1211	IFFI1212	IFFI1213	IFFI1214	IFFI1215
10	IFFI1220	IFFI1221	IFFI1224	IFFI1247	IFFI1248
	IFFI1251	IFFI1270	IFFI1277	IFFI1287	IFFI1289
	IFFI1290	IFFI1291	IFFI1292	IFFI1293	IFFI1297
	IFFI1300	IFFI1301	IFFI1302	IFFI1307	IFFI1308
	IFFI1309	IFFI1310	IFFI1311	IFFI1331	IFFI1335
15	IFFI1336	IFFI1337	IFFI1338	IFFI1339	IFFI1340
	IFFI1345	IFFI1348	IFFI1396	IFFI1397	IFFI1399
	IFFI1411	IFFI1413	IFFI1441	IFFI1443	
	<i>Saccharomyces cerevisiae</i> Hansen Var. ellipsoideus (Hansen) Dekker				
	ACCC2043	AS2.2	AS2.3	AS2.8	AS2.53
20	AS2.163	AS2.168	AS2.483	AS2.541	AS2.559
	AS2.606	AS2.607	AS2.611	AS2.612	
	<i>Saccharomyces chevalieri</i> Guilliermond				
	AS2.131	AS2.213			
	<i>Saccharomyces delbrueckii</i>				
25	AS2.285				
	<i>Saccharomyces delbrueckii</i> Lindner ver. mongolicus (Saito) Lodder et van Rij				
	AS2.209	AS2.1157			
	<i>Saccharomyces exiguous</i> Hansen				
	AS2.349	AS2.1158			
	<i>Saccharomyces fermentati</i> (Saito) Lodder et van Rij				
	AS2.286	AS2.343			
	<i>Saccharomyces logos</i> van laer et Denamur ex Jorgensen				
	AS2.156	AS2.327	AS2.335		

5	<i>Saccharomyces mellis</i> (Fabian et Quinet) Lodder et kreger van Rij				
	AS2.195				
	<i>Saccharomyces mellis</i> Microellipsoides Osterwalder				
	AS2.699				
	<i>Saccharomyces oviformis</i> Osteralder				
10	AS2.100				
	<i>Saccharomyces rosei</i> (Guilliermond) Lodder et Kreger van Rij				
	AS2.287				
	<i>Saccharomyces rouxii</i> Boutroux				
	AS2.178	AS2.180	AS2.370	AS2.371	
15	<i>Saccharomyces sake</i> Yabe				
	ACCC2045				
	<i>Candida arborea</i>				
	AS2.566				
	<i>Candida lambica</i> (Lindner et Genoud) van. Uden et Buckley				
20	AS2.1182				
	<i>Candida krusei</i> (Castellani) Berkhout				
	AS2.1045				
	<i>Candida lipolytica</i> (Harrison) Diddens et Lodder				
	AS2.1207	AS2.1216	AS2.1220	AS2.1379	AS2.1398
25	AS2.1399	AS2.1400			
	<i>Candida parapsilosis</i> (Ashford) Langeron et Talice Var. intermedia Van Rij et Verona				
	AS2.491				
	<i>Candida parapsilosis</i> (Ashford) Langeron et Talice				
	AS2.590				
	<i>Candida pulcherrima</i> (Lindner) Windisch				
	AS2.492				

<i>Candida rugousa</i> (Anderson) Diddens et Lodder				
AS2.511	AS2.1367	AS2.1369	AS2.1372	AS2.1373
AS2.1377	AS2.1378	AS2.1384		
<i>Candida tropicalis</i> (Castellani) Berkhout				
5	ACCC2004	ACCC2005	ACCC2006	AS2.164
	AS2.564	AS2.565	AS2.567	AS2.402
	AS2.637	AS2.1387	AS2.1397	AS2.568
				AS2.617
<i>Candida utilis</i> Henneberg Lodder et Kreger Van Rij				
	AS2.120	AS2.281	AS2.1180	
10	<i>Crebrothecium ashbyii</i> (Guilliermond)			
	Routein ( <i>Eremothecium ashbyii</i> Guilliermond)			
	AS2.481	AS2.482	AS2.1197	
<i>Geotrichum candidum</i> Link				
	ACCC2016	AS2.361	AS2.498	AS2.616
15	AS2.1062	AS2.1080	AS2.1132	AS2.1035
			AS2.1175	AS2.1183
<i>Hansenula anomala</i> (Hansen)H et P sydow				
	ACCC2018	AS2.294	AS2.295	AS2.296
	AS2.298	AS2.299	AS2.300	AS2.297
	AS2.339	AS2.340	AS2.341	AS2.302
20	AS2.641	AS2.642	AS2.782	AS2.338
			AS2.470	AS2.592
			AS2.635	AS2.794
<i>Hansenula arabitolgens</i> Fang				
	AS2.887			
<i>Hansenula jadinii</i> (A. et R Sartory Weill et Meyer) Wickerham				
	ACCC2019			
25	<i>Hansenula saturnus</i> (Klocker) H et P sydow			
	ACCC2020			
<i>Hansenula schneggii</i> (Weber ) Dekker				
	AS2.304			



<i>Hansenula subpelliculosa</i> Bedford				
AS2.740	AS2.760	AS2.761	AS2.770	AS2.783
AS2.790	AS2.798	AS2.866		
<i>Kloeckera apiculata</i> (Reess emend. Klocker) Janke				
5	ACCC2022	ACCC2023	AS2.197	AS2.496
	ACCC2021	AS2.711		AS2.714
<i>Lipomyces starkeyi</i> Lodder et van Rij				
	AS2.1390	ACCC2024		
<i>Pichia farinosa</i> (Lindner) Hansen				
10	ACCC2025	ACCC2026	AS2.86	AS2.87
	AS2.803			AS2.705
<i>Pichia membranaefaciens</i> Hansen				
	ACCC2027	AS2.89	AS2.661	AS2.1039
<i>Rhodospiridium toruloides</i> Banno				
15	ACCC2028			
<i>Rhodotorula glutinis</i> (Fresenius) Harrison				
	AS2.2029	AS2.280	ACCC2030	AS2.102
	AS2.278	AS2.499	AS2.694	AS2.107
	AS2.1146		AS2.703	AS2.704
20	<i>Rhodotorula minuta</i> (Saito) Harrison			
	AS2.277			
<i>Rhodotorula rubra</i> (Demme) Lodder				
	AS2.21	AS2.22	AS2.103	AS2.105
	AS2.140	AS2.166	AS2.167	AS2.108
25	AS2.282	ACCC2031		AS2.272
				AS2.279
<i>Rhodotorula aurantiaca</i> (Saito) Lodder				
	AS2.102	AS2.107	AS2.278	AS2.499
	AS2.703	AS2.1146		AS2.694

<i>Saccharomyces carlsbergensis</i> Hansen				
AS2.113	ACCC2032	ACCC2033	AS2.312	AS2.116
AS2.118	AS2.121	AS2.132	AS2.162	AS2.189
AS2.200	AS2.216	AS2.265	AS2.377	AS2.417
5 AS2.420	AS2.440	AS2.441	AS2.443	AS2.444
AS2.459	AS2.595	AS2.605	AS2.638	AS2.742
AS2.745	AS2.748	AS2.1042		
<i>Saccharomyces uvarum</i> Beijer				
IFFI1023	IFFI1032	IFFI1036	IFFI1044	IFFI1072
10 IFFI1205	IFFI1207			
<i>Saccharomyces willianus</i> Saccardo				
AS2.5	AS2.7	AS2.119	AS2.152	AS2.293
AS2.381	AS2.392	AS2.434	AS2.614	AS2.1189
<i>Saccharomyces sp.</i>				
15 AS2.311				
<i>Saccharomyces ludwigii</i> Hansen				
ACCC2044	AS2.243	AS2.508		
<i>Saccharomyces sinenses</i> Yue				
AS2.1395				
20	<i>Schizosaccharomyces octosporus</i> Beijerinck			
ACCC2046	AS2.1148			
<i>Schizosaccharomyces pombe</i> Lindner				
ACCC2047	ACCC2048	AS2.214	AS2.248	AS2.249
AS2.255	AS2.257	AS2.259	AS2.260	AS2.274
25 AS2.994	AS2.1043	AS2.1149	AS2.1178	IFFI1056
<i>Sporobolomyces roseus</i> Kluyver et van Niel				
ACCC2049	ACCC2050	AS2.19	AS2.962	AS2.1036
ACCC2051	AS2.261	AS2.262		

		<i>Torulopsis candida</i> (Saito) Lodder		
	AS2.270	ACCC2052		
		<i>Torulopsis famta</i> (Harrison) Lodder et van Rij		
	ACCC2053	AS2.685		
5		<i>Torulopsis globosa</i> (Olson et Hammer) Lodder et van Rij		
	ACCC2054	AS2.202		
		<i>Torulopsis inconspicua</i> Lodder et Kreger van Rij		
	AS2.75			
		<i>Trichosporon behrendii</i> Lodder et Kreger van Rij		
10	ACCC2056	AS2.1193		
		<i>Trichosporon capitatum</i> Diddens et Lodder		
	ACCC2056	AS2.1385		
		<i>Trichosporon cutaneum</i> (de Beurm et al.) Ota		
	ACCC2057	AS2.25	AS2.570	AS2.571 AS2.1374
15		<i>Wickerhamia fluorescens</i> (Soneda) Soneda		
	ACCC2058	AS2.1388		

## **II. Application of Electromagnetic Fields**

[0018] An electromagnetic field useful in this invention can be generated and applied by various means well known in the art. For instance, the EMF can be generated by applying an alternating electric field or an oscillating magnetic field.

[0019] Alternating electric fields can be applied to cell cultures through electrodes in direct contact with the culture medium, or through electromagnetic induction. See, e.g., Fig. 1. Relatively high electric fields in the medium can be generated using a method in which the electrodes are in contact with the medium.

Care must be taken to prevent electrolysis at the electrodes from introducing undesired ions into the culture and to prevent contact resistance, bubbles, or other features of electrolysis from dropping the field level below that intended. Electrodes should be matched to their environment, for example, using Ag-AgCl electrodes in solutions rich in chloride ions, and run at as low a voltage as possible.

For general review, see Goodman et al., *Effects of EMF on Molecules and Cells*, International Review of Cytology, A Survey of Cell Biology, Vol. 158, Academic Press, 1995.

5 [0020] The EMFs useful in this invention can also be generated by applying an oscillating magnetic field. An oscillating magnetic field can be generated by oscillating electric currents going through Helmholtz coils. Such a magnetic field in turn induces an electric field.

[0021] The frequencies of EMFs useful in this invention range from about 10200 to 13040 (e.g., 10200 to 10270, 12330 to 12390 and 12970 to 13040 MHz).  
10 Exemplary frequencies include 10231, 10237, 12361, 12997 and 13008 MHz. The field strength of the electric field useful in this invention ranges from about 20 to 600 mV/cm (e.g., 240-300, 310-340, 350-380, 380-430, 430-470 and 470-510 mV/cm). Exemplary field strengths include 246, 272, 288, 322, 343, 346, 364, 393, 446 and 483 mV/cm.

15 [0022] When a series of EMFs are applied to a yeast culture, the yeast culture can remain in the same container while the same set of EMF generator and emitters is used to change the frequency and/or field strength. The EMFs in the series can each have a different frequency or a different field strength; or a different frequency and a different field strength. Such frequencies and field strengths are  
20 preferably within the above-described ranges. Although any practical number of EMFs can be used in a series, it may be preferred that the yeast culture be exposed to a total of 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or more EMFs in a series. In one embodiment, the yeast culture is exposed to a series of EMFs, wherein the frequency of the electric field is alternated in the range of about 10200 to 10270,  
25 12330 to 12390 and 12970 to 13040 MHz.

[0023] Although the yeast cells can be activated after even a few hours of culturing in the presence of an EMF, it may be preferred that the compositions comprising activated yeast cells be allowed to multiply and grow in the presence of the EMF(s) for a total of about 140-210 hours. In the presence of 13 EMFs, the  
30 compositions can be grown for about 50-380, 80-380, 100-350, or 250-350 hours.

In the presence of 1 EMF, the compositions can be grown for about 5-60, 10-50, 80-180 or 100-150 hours.

[0024] Fig. 1 illustrates an exemplary apparatus for generating alternating electric fields. An electric field of a desired frequency and intensity can be  
5 generated by an AC source (3) capable of generating an alternating electric field, preferably in a sinusoidal wave form, in the frequency range of 5 to 20,000 MHz. Signal generators capable of generating signals with a narrower frequency range can also be used. If desired, a signal amplifier can also be used to increase the output. The culture container (2) can be made from a non-conductive material,  
10 e.g., glass, plastic or ceramic. The cable connecting the culture container (2) and the signal generator (3) is preferably a high frequency coaxial cable with a transmission frequency of at least 30 GHz.

[0025] The alternating electric field can be applied to the culture by a variety of means, including placing the yeast culture (1) in close proximity to the  
15 signal emitters such as a metal wire or tube capable of transmitting EMFs. The metal wire or tube can be made of red copper, and be placed inside the container (2), reaching as deep as 3-30 cm. For example, if the fluid in the container (2) has a depth of 15-20 cm, 20-30 cm, 30-50 cm, 50-70 cm, 70-100 cm, 100-150 cm or 150-200 cm, the metal wire can be 3-5 cm, 5-7 cm, 7-10 cm, 10-15 cm, 15-20 cm,  
20 20-30 cm and 25-30 cm from the bottom of the container (2), respectively. The number of metal wires/tubes used can be from 1 to 10 (e.g., 2 to 3). It is recommended, though not mandated, that for a culture having a volume up to 10 L, metal wires/tubes having a diameter of 0.5 to 2 mm be used. For a culture having a volume of 10-100 L, metal wires/tubes having a diameter of 3 to 5 mm can be  
25 used. For a culture having a volume of 100-1000 L, metal wires/tubes having a diameter of 6 to 15 mm can be used. For a culture having a volume greater than 1000 L, metal wires/tubes having a diameter of 20-25 mm can be used.

[0026] In one embodiment, the electric field is applied by electrodes submerged in the culture (1). In this embodiment, one of the electrodes can be a  
30 metal plate placed on the bottom of the container (2), and the other electrode can comprise a plurality of electrode wires evenly distributed in the culture (1) so as to

achieve even distribution of the electric field energy. The number of electrode wires used depends on the volume of the culture as well as the diameter of the wires.

### **III. Culture Media**

5 [0027] Culture media useful in this invention contain sources of nutrients assimilatable by yeast cells. Complex carbon-containing substances in a suitable form (e.g., carbohydrates such as sucrose, glucose, dextrose, maltose, starch and xylose; or mannitol) can be the carbon sources for yeast cells. The exact quantity of the carbon sources can be adjusted in accordance with the other ingredients of  
10 the medium. In general, the amount of carbon-containing substances varies between about 0.5% and 10% by weight of the medium, and preferably between about 1% and 5%, most preferably between about 1.0-2.0%. Vitamins can also be added to the medium, for example, Vitamin E, D<sub>3</sub>, H and B<sub>6</sub>. Among the inorganic salts which can be added to a laboratory culture medium are the  
15 customary salts capable of yielding sodium, potassium, calcium, phosphate, sulfate, carbonate, and like ions. Non-limiting examples of nutrient inorganic salts are (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub>, CaCO<sub>3</sub>, KH<sub>2</sub>PO<sub>4</sub>, K<sub>2</sub>HPO<sub>4</sub>, MgSO<sub>4</sub>, NaCl, and CaSO<sub>4</sub>.

### **IV. Electromagnetic Activation of Yeast Cells**

[0028] To activate or enhance the innate ability of yeast cells to produce  
20 agents that are useful in regulating the central nervous system, these cells can be cultured in an appropriate medium under sterile conditions at 20°C-35°C (e.g., 28-32°C) for a sufficient amount of time, e.g. 5-60, 10-50, 80-180, 100-150, 140-210, 50-380, 80-380, 100-350, or 250-350 hours in an alternating electric field or a series of alternating electric fields as described above.

25 [0029] An exemplary set-up of the culture process is depicted in Fig. 1 (see above). An exemplary culture medium contains the following in per 1000 ml of sterile water: 6.0 g of sucrose, 12 g of mannitol, 60 µg of Vitamin E, 50 µg of Vitamin D<sub>3</sub>, 60 µg Vitamin H, 90 µg of Vitamin B<sub>6</sub>, 50 ml bovine serum, 0.2 g of KH<sub>2</sub>PO<sub>4</sub>, 0.25 g of MgSO<sub>4</sub>•7H<sub>2</sub>O, 0.3 g of NaCl, 0.2 g of CaSO<sub>4</sub>•2H<sub>2</sub>O, 4.0 g of

CaCO<sub>3</sub>•5H<sub>2</sub>O and 2.5 g of peptone. All vitamins are sterilized before added to the solution. Yeast cells of the desired strains are then added to the culture medium to form a mixture containing  $1 \times 10^8$  yeast cells per 1000 ml of culture medium. The yeast cells can be of any of the strains illustrated in Table 1. In one embodiment, 5 the yeast cells are of the strain *Saccharomyces cerevisiae* Hansen IFFI1335. The mixture is then added to the apparatus of Fig. 1.

[0030] The activation process of the yeast cells involves the following steps: 1) maintaining the temperature of the activation apparatus at 20-35°C (e.g., 28-32°C), and culturing the yeast cells for 28 hours; 2) applying an electric field 10 having a frequency of about 10231 MHz and a field strength of 240-260 mV/cm (e.g., about 246 mV/cm) for 16 hours; 3) then applying an electric field having a frequency of about 10237 MHz and a field strength of 310-340 mV/cm (e.g., about 322 mV/cm) for 42 hours; 4) then applying an electric field having a frequency of about 12361 MHz and a field strength of 350-380 mV/cm (about 364 mV/cm) for 15 38 hours; 5) then applying an electric field having a frequency of about 12997 MHz and a field strength of 380-420 mV/cm (e.g., about 393 mV/cm) for 38 hours; 6) then applying an electric field having a frequency of about 13008 MHz and a field strength of 280-300 mV/cm (e.g., about 288 mV/cm) for 16 hours; and 7) finally lyophilizing the activated yeast cells to form a powder and storing the 20 powder at 4 °C. Preferably, the concentration of the lyophilized yeast cells is more than  $10^{10}$  cells/g.

#### **V. Acclimatization of Yeast Cells To the Gastric Environment**

[0031] Because the yeast compositions of this invention must pass through the stomach before reaching the small intestine, where the effective components 25 are released from these yeast cells, it is preferred that these yeast cells be cultured under acidic conditions to acclimatize the cells to the gastric juice. This acclimatization process results in better viability of the yeasts in the acidic gastric environment.

[0032] To achieve this, the yeast powder containing activated yeast cells 30 can be mixed with an acclimatizing culture medium at 10 g (containing more than

- 10<sup>10</sup> activated cells per gram) per 1000 ml. The yeast mixture is then cultured first in the presence of an alternating electric field having a frequency of about 12997 MHz and a field strength of 370-430 mV/cm (e.g., about 446 mV/cm) at about 28 to 32°C for 34-42 hours (e.g., 38 hours). The resultant yeast cells are further
- 5 incubated in the presence of an alternating electric field having a frequency of about 13008 MHz and a field strength of 350-380 mV/cm (e.g., about 364 mV/cm) at about 28 to 32°C for 16-28 hours (e.g., 20 hours). The resulting acclimatized yeast cells are then dried and stored either in powder form ( $\geq 10^{10}$  cells/g) at room temperature or stored in vacuum at 0-4°C.
- 10 [0033] An exemplary acclimatizing culture medium is made by mixing 700 ml of fresh pig gastric juice and 300 ml of wild Chinese hawthorn extract. The pH of acclimatizing culture medium is adjusted to 2.5 with 0.1 M hydrochloric acid and 0.2 M potassium biphthalate ( $C_6H_4(COOK)COOH$ ). The fresh pig gastric juice is prepared as follows. At about 4 months of age, newborn Holland white
- 15 pigs are sacrificed, and the entire contents of their stomachs are retrieved and mixed with 2000 ml of water under sterile conditions. The mixture is then allowed to stand for 6 hours at 4°C under sterile conditions to precipitate food debris. To prepare the wild Chinese hawthorn extract, 500 g of fresh wild Chinese hawthorn is dried under sterile conditions to reduce the water content ( $\leq 8\%$ ). The dried fruit
- 20 is then ground ( $\geq 20$  mesh) and added to 1500 ml of sterile water. The mixture is allowed to stand for 6 hours at 4°C under sterile conditions. The supernatant is collected to be used in the acclimatizing culture medium.

## **VI. Manufacture of Yeast Compositions**

- [0034] To prepare the yeast compositions of the invention, an apparatus
- 25 depicted in Fig. 2 or an equivalent thereof can be used. This apparatus includes a first container (A), a second container (B), and a third container (C), each equipped with a pair of electrodes (4). One of the electrodes is a metal plate placed on the bottom of the containers, and the other electrode comprises a plurality of electrode wires evenly distributed in the space within the container to achieve even



distribution of the electric field energy. All three pairs of electrodes are connected to a common signal generator.

[0035] The culture medium used for this purpose is a mixed fruit extract solution containing the following ingredients per 1000 L: 300 L of wild Chinese hawthorn extract, 300 L of jujube extract, 300 L of fruit extract from *Schisandra chinensis* Baill (wu wei zi), and 100 L of soy bean extracts. To prepare hawthorn, jujube and wu wei zi extracts, the fresh fruits are washed and dried under sterile conditions to reduce the water content to no higher than 8%. One hundred kilograms of the dried fruits are then ground ( $\geq 20$  mesh) and added to 400 L of sterile water. The mixtures are stirred under sterile conditions at room temperature for twelve hours, and then centrifuged at 1000 rpm to remove insoluble residues. To make the soy bean extract, fresh soy beans are washed and dried under sterile conditions to reduce the water content to no higher than 8%. Thirty kilograms of dried soy beans are then ground into particles of no smaller than 20 mesh, and added to 130 L of sterile water. The mixture is stirred under sterile conditions at room temperature for twelve hours and centrifuged at 1000 rpm to remove insoluble residues. Once the mixed fruit extract solution is prepared, the solution is sterilized at 121°C for 30 minutes, and cooled to 40°C before use.

[0036] One thousand grams of the activated yeast powder prepared as described above (Section V, *supra*) is added to 1000 L of the mixed fruit extract solution, and the yeast solution is transferred to container (A) shown in Fig. 2. The yeast cells are then cultured in the presence of an alternating electric field having a frequency of about 12997 MHz and a field strength of about 430-470 mV/cm (e.g., about 446 mV/cm) at 28-32°C under sterile conditions for 38 hours. The yeast cells are further incubated in an alternating electric field having a frequency of about 13008 MHz and a field strength of 330-360 mV/cm (e.g., about 343 mV/cm). The culturing continues for another 12 hours.

[0037] The yeast culture is then transferred from the first container (A) to the second container (B) (if need be, a new batch of yeast culture can be started in the now available first container (A)), and subjected to an alternating electric field having a frequency of about 12997 MHz and a field strength of 470-510 mV/cm

(e.g., about 483 mV/cm) for 24 hours. Subsequently the frequency and field strength of the electric field are changed to about 13008 MHz and 350-380 mV/cm (e.g., about 368 mV/cm), respectively. The culturing continues for another 12 hours.

5 [0038] The yeast culture is then transferred from the second container (B) to the third container (C), and subjected to an alternating electric field having a frequency of about 12997 MHz and a field strength of 330-360 mV/cm (e.g., about 346 mV/cm) for 28 hours. Subsequently the frequency and field strength of the electric field are changed to about 13008 MHz and 260-280 mV/cm (e.g., about  
10 272 mV/cm), respectively. The culturing continues for another 12 hours.

[0039] The yeast culture from the third container (C) can then be packaged into vacuum sealed bottles of 30-50 ml or 100 ml for use as a dietary supplement, e.g., health drinks, or medication in the form of pills, powder, etc. The dietary supplement can be taken 3-4 times daily at 30-60 ml each time for a period of three  
15 months (10-30 minutes before meals and at bedtime). If desired, the final yeast culture can also be dried within 24 hours and stored in powder form.

[0040] In one embodiment, the compositions of the invention can also be administered intravenously or peritoneally in the form of a sterile injectable preparation. Such a sterile preparation is prepared as follows. A sterilized health  
20 drink composition is first treated under ultrasound ( $\geq 18,000$  Hz) for 10 minutes and then centrifuged at 4355 rpm for another 10 minutes. The resulting supernatant is adjusted to pH 7.2-7.4 using 1 M NaOH and subsequently filtered through a membrane (0.22  $\mu\text{m}$  for intravenous injection and 0.45  $\mu\text{m}$  for peritoneal injection) under sterile conditions. The resulting sterile preparation is submerged  
25 in a 35-38 °C water bath for 30 minutes before use.

[0041] The yeast compositions of the present invention are derived from yeasts used in food and pharmaceutical industries. The yeast compositions are thus devoid of side effects associated with many pharmaceutical compounds.

[0042] In other embodiments, the compositions of the invention may also be  
30 formulated with pharmaceutically acceptable carriers to be orally administered in

any orally acceptable dosage form including, but not limited to, capsules, tablets, suspensions or solutions.

### Examples

[0043] The following examples are meant to illustrate the methods and materials of the present invention. Suitable modifications and adaptations of the described conditions and parameters which are obvious to those skilled in the art are within the spirit and scope of the present invention.

[0044] The activated yeast compositions used in the following experiments were prepared as described above, using *Saccharomyces cerevisiae* Hansen IFFI1335 cultured in the presence of an alternating electric field having the electric field frequency and field strength exemplified in the parentheses following the recommended ranges in Section IV, *supra*. Control yeast compositions were those prepared in the same manner except that the yeast cells were cultured in the absence of EMFs. Unless otherwise indicated, the yeast compositions and the corresponding controls were admitted to the animals via intragastric feeding.

#### Example 1: Anti-seizure Effect on Electroshocked Mice

[0045] Electrodes are placed at the ears of the mice to provide instant stimulation through a strong electric current. An appropriate electric current induces an excessive surge of electrical activity in the brain resulting in seizures to occur.

[0046] Kunming grade healthy mice, which were 50-70 days old, weighing 18-22 g, were provided by the Institute of Zoology, Chinese Academy of Sciences, Beijing, China. The electroshock device (model GJ-2, manufactured by Medical Instrument Factory, Zhejiang Medical University) was adjusted to an output frequency of 60 Hz and voltage of 125 V. An equal number of male and female mice were selected. The ears of the mice were covered with saturated saline soaked-cotton. The electrodes were then clipped onto the ears of the mice. The mice were electroshocked three times for 0.3 seconds at an interval of 10 seconds. Mice with seizure activity were then divided into four groups, each group

containing 70 mice: the test group (AY), the control yeast group (NY), the positive control group (CK2) and negative control group (CK1). Each mouse in the test, control yeast and negative control groups was administered twice daily 0.4 ml of the activated yeast composition, the control yeast composition, and saline, respectively, for 1 week. Each mouse in the positive control group was administered twice daily 15 mg/kg of phenobarbital for 1 week.

[0047] On each day, the mice were electroshocked once for 0.3 seconds.

The number of mice with seizure activity were monitored as illustrated in Table 2.

Table 2

Group	Day 1		Day 2		Day 3		Day 4	
	No. mice with seizure activity	% mice with anti-seizure	No. mice with seizure activity	% mice anti-seizure	No. mice with seizure activity	% mice anti-seizure	No. mice with seizure activity	% mice anti-seizure
AY	67	4.3	56	20	44	37.1	32	54.3
NY	70	0	70	0	70	0	69	1.4
CK2	59	15.7	48	31.4	56	20	55	21.4
CK1	70	0	70	0	70	0	70	0

Group	Day 5		Day 6		Day 7	
	No. mice with seizure activity	% mice anti-seizure	No. mice with seizure activity	% mice anti-seizure	No. mice with seizure activity	% mice anti-seizure
AY	17	75.7	4	94.3	1	98.6
NY	70	0	70	0	70	0
CK2	58	17.1	61	12.8	61	12.9
CK1	70	0	70	0	70	0

[0048] As illustrated above, compared to the control (CK1) and control yeast (NY) groups, the test group (AY) shows a significant decrease in the number of rats with seizure activity after day 4. Further, nearly all of the rats in the test group show an anti-seizure effect after day 7, while only 12.9% of the rats in the

positive control group (CK2) have an anti-seizure effect . Thus, the activated yeast composition of this invention has an anti-seizure affect on electro-shocked mice.

Example 2: Anti-seizure effect of rats induced with cardiazol

- [0049] Cardiazol induces seizure activity through intensifying excitatory synaptic activity of the brain. At the appropriate dose, cardiazol induces epileptic bursts that occur spontaneously.
- [0050] An equal number of female and male Wistar rats that were 5-6 months old, weighing 200-220 g were provided by the Chinese Academy of Military Medical Sciences, Beijing, China. The rats were divided into four groups, each containing 20 rats: the test group (AY), the control yeast group (NY), the positive control group (CK2) and negative control group (CK1). Each rat in the test, control yeast and negative control groups was administered twice daily 0.6 ml of the activated yeast composition, the control yeast composition, and saline, respectively, for 1 week. Each mouse in the positive group was administered twice daily 0.04 g/kg (body weight) of phenobarbital (dissolved in 0.6 ml saline) for 1 week.
- [0051] On day 7, 30 minutes after administering the last dosage of the above compositions, each group was administered by subcutaneous injection 70 mg/kg (body weight) of cardiazol (prepared as a 5% cardiazol solution in saline). The number of seizures occurred, incubation time (time between completion of injection and first seizure) and life span after injection was recorded immediately thereafter as shown in Table 3.

Table 3

Group	Animal number	incubation time (min)	life span (min)
AY	20	11.82 ± 1.52	68.33 ± 16.26
NY	20	2.52 ± 1.16	8.23 ± 3.42
CK2	20	7.23 ± 5.32	22.76 ± 17.37

CK1	20	$2.60 \pm 1.21$	$7.62 \pm 3.65$
-----	----	-----------------	-----------------

[0052] As illustrated above, compared to the control yeast (NY), negative control groups (CK1), the test group (AY) shows a significant increase in the incubation time of the seizure and the life span of the rat after cardiazol injection.

5 Further, compared to the positive control group (CK2), the test group shows that the incubation time of the seizure as well as the life span of the rat after cardiazol injection is much higher. Thus, the activated composition of this invention has an anti-seizure effect on rats treated with a seizure-inducing drug, cardiazol.

[0053] While a number of embodiments of this invention have been set  
10 forth, it is apparent that the basic constructions may be altered to provide other embodiments which utilize the compositions and methods of this invention.